

## **KANGAROO CARE – NICU ANTIBIOTIC POLICY**

Infection has become an increasingly important cause of mortality and morbidity in NICUs. Infection usually falls into two main categories: early onset and late onset sepsis. The cut off between early and late onset is usually taken as 48-72 hr.

### **Risk Factors:**

#### **Early onset of Sepsis:**

1. UTI in pregnancy
2. Preterm labour
3. Premature rupture of membranes
4. Prolonged rupture of membranes (>18 hours)
5. Maternal fever
6. Maternal carrier Group B Streptococcus
7. Maternal chorioamnionitis
8. Outborn neonates

#### **Organisms responsible for early onset sepsis:**

The predominant pathogens come from the maternal genital tract and are:

- E.coli
- Klebsiella spp
- Pseudomonas spp
- Group B Streptococcus (GBS)

#### **Late onset of Sepsis:**


1. Deficient immune responses in ELBW infants
2. Central lines, especially UVC > 10 days
3. Outborn neonates
4. Colonization of newborns by MRSA
5. Inadequate infection control practices by staff and visitors.
6. Pre mature neonates

#### **Organisms responsible for late onset sepsis:**

The predominant pathogens are:

- Gram Positive organisms
  - Coagulase- negative staphylococci
  - Staphylococcus aureus

- Enterococci
- Streptococcus spp
- Gram Negative organisms
  - E coli
  - Klebsiella spp
  - Pseudomonas spp
  - Acinetobacter spp



- \* *Full diagnostic evaluation includes a blood culture, a complete blood count (CBC) including white blood cells, differential and platelet counts, chest radiography (if respiratory abnormalities are present), and lumbar puncture (if patient is stable enough to tolerate procedure and sepsis is suspected).*
- ~ *Antibiotic therapy should be directed toward the most common causes of neonatal sepsis, including intravenous ampicillin for GBS and coverage for other organisms (including Escheria.coli and other gram – negative pathogens) and should take into account local antibiotic resistance patterns.*
- \* *Consultations with obstetric providers is important to determine the level of clinical suspicious for chorioamnionitis, chorioamnionitis is diagnosed clinically and some of the signs are no specific.*

### Early onset sepsis

#### Baby < 28 weeks

- Without risk factors: *IV Ampicillin and Gentamycin (Consider Amikacin for out-born babies instead of Gentamycin) for 5 days*
- With risk factors: *Cefotaxime and Amikacin for 5-7 days*
- Second line: *IV Piptaz and Amikacin for 5-7 days*

#### Babies 28-32 weeks:

- Without risk factors: *Amoxicillin-Clavulanic Acid and Gentamycin (Consider Amikacin for outborn babies instead of Gentamycin) for 5 days*
- With risk factors: *Cefotaxime and Gentamycin for 5-7 days*  
Second line of antibiotics: *IV Piptaz and Amikacin for 5-7 days*

#### Babies > 32 weeks:

- Without risk factors: *Amoxicillin-Clavulanic Acid and Gentamycin (Consider Amikacin for outborn babies instead of Gentamycin) for 5 days*
- With risk factors: *Cefotaxime and Gentamycin for 5-7 days*  
Second Line of Antibiotics: *Imipenem and Amikacin for 5-7 days*

**(Considered Imipenem if meningitis not suspected and Meropenem for suspected meningitis)**

**Add metronidazole if anaerobic infection is suspected.**

Less than 28 weeks		
	No risk factors	Ampicillin and Gentamycin
	With risk factors	Cefotaxime and Amikacin
	Second line	Piptaz and Amikacin
28-32 weeks		
	No risk factors	Augmentin and Gentamycin
	With risk factors	Cefotaxime and Gentamycin
	Second line	Piptaz and Amikacin
More than 32 weeks		
	No risk factors	Augmentin and Gentamycin
	With risk factors	Cefotaxime and Gentamycin
	Second line	Imipenem/Meropenem and Amikacin

### Late onset sepsis

**IV Amoxicillin-Clavulanic Acid + Amikacin** till we get PCR or blood culture report.

Second Line of Antibiotics: **Cefaperazone-Sulbactam and Amikacin**

- **Ceftriaxone** can be considered in place of Cefotaxime in term infants.
- Add **metronidazole** if suspicion of anaerobic infection (e.g. intra-abdominal sepsis, NEC)
- IV **Vancomycin** for Gram positive sepsis, and in MRSA
- Consider **Piperacillin- Tazobactam and Amikacin** for ventilator-associated pneumonia
- Duration of antibiotics depends on risk factors, clinical condition and lab reports; kindly review antibiotics after 48-72 hrs
- Incubator will be used till 28 weeks only and will be changed every week  Incubator humidifier will be changed every 72 hrs and sent to ETO.

**Fluconazole will be given if 2 out of the following conditions are met**

- Birth weight less than 1kg.
- TPN
- On Ventilator
- Broad Spectrum antibiotics
- Central line

### **Amphotericin B**

- In proven fungal infections, liposomal form of the same should be given

## Second Line

- 5 Flucytosine, kindly do CBC before starting. Flucytosine is given along with Amphotericin B in CNS infections

Review Date	February 2025
Next Review Date	February 2026
Approval	KC Neonatal Service Clinical Guideline Meeting
Authors	Dr Shekar Subbaiah / Dr Ramapriya Kalkunte/ Dr Kishore Yerur / Dr Shalmali
Patient to whom this applies	Patients of the KC Neonatal Service and newborn infants on the Postnatal Wards and Labour Suites of the KC Hospitals who fit the inclusion criteria of the guideline
Evidence used	The contemporary evidence base has been used to develop this guideline.